

## Mitophagy Pathways in Health and Disease

Peripartum cardiomyopathy (PPCM) is a relatively rare form of non-ischemic cardiomyopathy that occurs in healthy women during the final month of pregnancy and up to 5 months post-delivery. According to the National Institutes of Health consensus panel, diagnostic criteria for PPCM includes congestive heart failure symptoms of unknown cause, absence of a pre-existing heart muscle disorder, symptomatic onset during the last month of gestation or 5 months postpartum, and left ventricular systolic dysfunction demonstrated by a decreased ejection fraction (EF) (Pearson et al., 2000). Its incidence in United States varies from 1 in 1300 to 1 in 4000 live births. (Message from the Guest Editors)

Peripartum cardiomyopathy (PPCM) is a relatively rare form of non-ischemic cardiomyopathy that occurs in healthy women during the final month of pregnancy and up to 5 months post-delivery. According to the National Institutes of Health consensus panel, diagnostic criteria for PPCM includes congestive heart failure symptoms of unknown cause, absence of a pre-existing heart muscle disorder, symptomatic onset during the last month of gestation or 5 months postpartum, and left ventricular systolic dysfunction demonstrated by a decreased ejection fraction (EF) (Pearson et al., 2000). Its incidence in United States varies from 1 in 1300 to 1 in 4000 live births. (Message from the Guest Editors)

## Guest Editors:

### Roman R. Poznanski

Universiti Sultan Zainal Abidin, Malaysia and Rockefeller University. Peripartum cardiomyopathy (PPCM) is a relatively rare form of non-ischemic cardiomyopathy that occurs in healthy women during the final month of pregnancy and up to 5 months post-delivery (Johnson-Coyle et al., 2012).



**Roman R. Poznanski**

**Submission Deadline:** 31 Dec 2020

**Submission:** JIN special issue

**Contact us:** jin



**IMR PRESS**